



## RESEARCH PAPER

## OPEN ACCESS

## Heritability and genetic diversity of iron, zinc and some morphological and physiological traits in some spring wheat genotypes (*Triticum aestivum* L.)

Mostafa Khodadadi<sup>1</sup>, Hamid Dehghani<sup>1\*</sup>, Mohammad Hussein Fotokian<sup>2</sup>

<sup>1</sup>Plant Breeding and Biotechnology Department, Faculty of Agriculture, Tarbiat Modares University, Tehran, Iran

<sup>2</sup>Plant Breeding and Agronomy Department, Faculty of Agriculture, Shahed University, Tehran, Iran

**Key words:** Clustering, iron, wheat, zinc.

doi: <http://dx.doi.org/10.12692/ijb/4.3.139-148>

Article published on February 14, 2014

### Abstract

Wheat is one of the most important cereals and main source of food in Iran and many different countries. Thirty spring common wheat genotypes from Iran were assessed for grain concentrations of Iron and Zinc and some morphological and physiological traits in 2009 and 2010. Iron showed large variation among genotypes, ranging from 17.5  $\mu\text{g g}^{-1}$  to 48.68  $\mu\text{g g}^{-1}$  (mean 30.79  $\mu\text{g g}^{-1}$ ) in 2009 and from 22.74  $\mu\text{g g}^{-1}$  to 45.60  $\mu\text{g g}^{-1}$  (mean 30.53  $\mu\text{g g}^{-1}$ ) in 2010. Similarly, Zinc concentration varied among genotypes, ranging from 22.36  $\mu\text{g g}^{-1}$  to 52.69  $\mu\text{g g}^{-1}$  (mean 35.79  $\mu\text{g g}^{-1}$ ) in 2009 and from 20.74  $\mu\text{g g}^{-1}$  to 78.60  $\mu\text{g g}^{-1}$  (mean 42.45  $\mu\text{g g}^{-1}$ ) in 2010. The highest heritability was exhibited by days to 50% heading (0.96) in 2009 and by stem number (0.96), spike length (0.96) in 2010. Heritability of grain iron and zinc were 0.74 and 0.61 in 2009, 0.85 and 0.92 in 2010 respectively. Four clusters formed through cluster analysis. Sistan, Akbari, Roshan and Roshan Bahareh backcross genotypes are recommended to use in multiple crossing program to reach transgressive segregants with high genetic potential of studied traits especially for chlorophyll content, grain iron and grain zinc all together. The most squared Euclidean genetic distance (75.81) observed between Niknejad and Roshan genotypes. Therefore, these genotypes can be used to plan wide crosses, to reach genetic diversity and maximize expression of heterosis. Mahdavi and Bahar genotypes had the most squared Euclidean genetic distance (556.45) for grain iron and zinc content.

\* Corresponding Author: Hamid Dehghani ✉ [dehghanirh@yahoo.com](mailto:dehghanirh@yahoo.com)

## Introduction

Approximately 600 million metric tons of wheat are commercially milled annually and consumed in nearly every nation of the world (Akhtar and Ashgar, 2011). Whole wheat flour is a staple food of the Indian population and supply more than 50% of the total energy intake. Similarly, in countries such as Syria, Algeria, Turkey, and Iran, wheat consumption is approximately half the total caloric intake and ranges up to 600 g/day (Ranum, 2001). Zinc and iron are essential micronutrients for human nutrition and health. There are estimates that some three billion people worldwide are afflicted by iron deficiency, and up to half of the population in developing countries are at risk of zinc deficiency (Welch and Graham, 2004; Cakmak, 2008). Micronutrient malnutrition greatly increases mortality and morbidity rates, diminishes cognitive abilities of children and lowers their educational attainment, reduces labor productivity, stagnates national development efforts, contributes to continued high population growth rates and reduces the livelihood and quality of life for all those affected (Welch and Graham, 2002). Zinc and iron deficiencies are particularly known to be a common problem in populations that depend on cereals as the main staple food and with little or no access to animal products (White and Broadley, 2005). According to above mentioned subjects, wheat flour and its products are good cases for fortification with iron and zinc. But, fortification efforts are highly dependent on funding, and the scope is restricted to a single geographical area. Standard fortification programs must be sustained at the same level of funding year after year; and if the investments are not sustained, the benefits disappear, in contrast micronutrient improvement by plant breeding programs is sustainable successful (Bouis *et al.*, 2000). Consequently, genetically improvement of wheat grain Iron and Zinc concentration is a good approach. Previously, Morganov *et al.* (2007) studied on sixty-six spring and winter wheat cultivars and advanced lines from Central Asian national breeding programs of the Central Asia region comprises five countries (Kazakhstan, Kyrgyzstan, Turkmenistan,

Tajikistan and Uzbekistan) to determine the levels of Iron and Zinc in the grain of the target wheat genotypes and identify promising lines with higher Iron and Zinc concentrations in the grain. Also, cakmak *et al.* (2004) studied on 825 wild emmer accessions (*Triticum turgidum* ssp. *dicoccoides*) for natural variation of Iron and Zinc as an important genetic resource for improving zinc and iron concentration in cultivated wheat.

One of the important goals to wheat breeding is hybridization and subsequent selection. Thus select of parents with the most genetic distance is the first step in breeding program. However, genetic distance between parents is necessary to develop superior hybrid (Ilker *et al.*, 2010). Transgressive segregation occurs often when the difference between parents is small and the additive variance is high (Joshi *et al.*, 2004) and it is better that these parents selected from divergent cluster with high performance for under breeding traits (Kumar *et al.*, 2009).

In order to determine whether iron and zinc concentration in a specific crop or other traits can be improved by traditional breeding methods, it must be known to how much these traits are heritable (Courtney, 2007). Heritability is a measure of the extent to which observed phenotypic differences for a trait are due to genetic variations (Klug and Cummings, 2005). A study of genetic variability with the help of suitable parameters such as genotypic coefficient of variation and heritability are thoroughly necessary to start an efficient breeding program (Mishra *et al.*, 1988). Although direct selection for various traits could be misleading, indirect selection via related traits with high heritability might be more impressive than direct selection (Toker and Cigirgan, 2004). Targeted opt of physiological traits that limit yield and have a high heritability may be more impressive than direct selection for yield (Sayar *et al.*, 2007). Iron and zinc concentration, exists in the germplasm for the crop. Thus, an important initial step in developing a biofortified crop is the screening of germplasm to see if this genetic variation exists.

Such germplasm screenings have been undertaken for a number of crops. Studies with common bean (*Phaseolus vulgaris*), rice (*Oriza sativa*), and wheat (*Triticum spp.*) have all found large variation in iron and zinc concentration (Welch and Graham, 2004).

The genetic variation of selected parents is not always based on factors such as geographic diversity place of release of ploidy level. So, characterization of genotypes should be based on statistical procedure such as hierarchical Euclidean cluster analysis (Kumar *et al.*, 2009). The objectives of this study was determine the level of grain iron and zinc content in the Iranian spring wheat genotypes and identify genotypes with the highest genetic distance for using in breeding programs.

### Material and methods

#### *Plant materials and experimental design*

Thirty spring common wheat genotypes (Fig. 1) supplied by the Agricultural Research Institutes gene bank in Karaj were prepared and used for this research in the Research Field of Faculty of Agricultural Sciences, Tarbiat Modares University, Tehran, Iran in 2009 and 2010. The plot size was 1.5 m length with three rows, row spacing was 20 cm and seeding rate was 250 seeds m<sup>-2</sup> based on a randomized complete block design with three replications. Nitrogen and phosphorus fertilizer, were consumed at 40 and 60 Kg ha<sup>-1</sup>, respectively before planting and nitrogen fertilizer was also broadcast at the stages of tillering and stem elongation at 40 kg ha<sup>-1</sup>. The measured traits include stem number, fertile stem number, flag leaf length (cm), flag leaf width (cm), leaf area (cm<sup>2</sup>), main stem leaf number, spike length (cm), days to 50% heading, 1000 grain weight, chlorophyll content (measured by SPAD-502), grain iron concentration (µg g<sup>-1</sup>) and grain zinc concentration (µg g<sup>-1</sup>).

Grain iron and zinc were measured using atomic absorption method (Gupta, 2000) after digestion based on nitric (0.1 normal)/citric (10%) acid (Morgounov *et al.* 2007) and extraction by ion-chromatography method (Small *et al.*, 1975). Also,

homogeneity of experimental field has confirmed and then trial was conducted.

#### *Statistical analysis*

Normality test using Shapiro-Wilk method, analysis of variance, means comparison by Duncan's new multiple range test (DMRT) at 5% level of probability and genotypes grouping using hierarchical Euclidean cluster analysis (Kumar *et al.*, 2009) were performed using the statistical software SPSS version 16.0 (SPSS, Chicago, USA) program. In addition, cophenetic correlation computed through Eta statistic using SPSS. For cluster analysis, ward's method based on squared Euclidean distance measure was used. Because of non-uniformity of measurement scale of traits data were standardized (Mohammadi and prasanna, 2003) using Z score method while in cluster analysis for grain iron and zinc original data were used. Some statistics including squared distance between cluster centroids, within cluster sum of square, average and maximum distance from centroid computed using Minitab 14.

The mean squares were used to estimate genotypic and phenotypic variance according to Johnson *et al.* (1955). The coefficient of variation was calculated according to the formula suggested by Burton (1952). The phenotypic and genotypic coefficient of variation and broadcast heritability were calculated according to Singh and Chowdhury (1985).

### Results

#### *Analysis of variance for studied traits*

Results of normality test showed that data for all traits were normal except for grain iron and zinc in second year. Therefore, data transformation for these traits was done. Analysis of variance showed high significant difference among genotypes for all traits in both years (Table 1). Some descriptive statistics of studied traits were presented in Table 3. Sistan, Roshan Bahareh backcross, Chenab, Bayat, Kavir, Arta, Shiraz, Shiraz and Bayat, Dez, Verinak, Akbari and Akbari genotypes in 2009; Sholeh, Darya, Sistan, Roshan-Bahareh back cross, Roshan, Roshan and Bayat, Shiraz, Shiraz and Bayat, Akbari, Roshan, Marvdasht and Darab 2 genotypes in 2010 exhibited

maximum means for traits presented in Table 3 with order from left to right.

**Table 1.** Analysis of variance for studied traits in 30 spring wheat genotypes.

Year	S.O.V	DF	Mean square											
			Stem No.	Fertile stem No.	Flag leaf length	Flag leaf width	Leaf area	Main stem leaf No.	Spike length	Days to 50% heading	1000 grain weight	Chlorophyll content	Grain iron	Grain zinc
2009	Replication	2	18.00 <sup>*</sup>	4.00 <sup>ns</sup>	27.00 <sup>**</sup>	0.02 <sup>ns</sup>	5.00 <sup>ns</sup>	1.00 <sup>*</sup>	0.35 <sup>ns</sup>	2.00 <sup>ns</sup>	8.98 <sup>*</sup>	138 <sup>***</sup>	145.21 <sup>**</sup>	0.47 <sup>ns</sup>
	Genotype	29	16.3 <sup>***</sup>	4.00 <sup>***</sup>	16.20 <sup>***</sup>	0.08 <sup>***</sup>	35.7 <sup>**</sup>	0.6 <sup>***</sup>	2.3 <sup>***</sup>	19.90 <sup>***</sup>	28.78 <sup>***</sup>	32.1 <sup>***</sup>	89.71 <sup>***</sup>	61.90 <sup>***</sup>
	Error	58	5.10	1.70	5.80	0.02	15.90	0.22	0.29	0.70	2.73	8.40	23.07	23.90
2010	Replication	2	0.27 <sup>ns</sup>	3.74 <sup>***</sup>	68.21 <sup>***</sup>	0.62 <sup>***</sup>	9.93 <sup>ns</sup>	0.08 <sup>ns</sup>	2.58 <sup>***</sup>	4.58 <sup>*</sup>	1.14 <sup>ns</sup>	5.43 <sup>ns</sup>	1.73e <sup>-6</sup> <sup>ns</sup>	0.002 <sup>ns</sup>
	Genotype	29	8.52 <sup>***</sup>	4.78 <sup>***</sup>	28.86 <sup>***</sup>	0.07 <sup>***</sup>	48.17 <sup>***</sup>	0.31 <sup>***</sup>	3.11 <sup>***</sup>	27.39 <sup>***</sup>	64.02 <sup>***</sup>	32.81 <sup>***</sup>	6.71e <sup>-5</sup> <sup>***</sup>	0.05 <sup>***</sup>
	Error	58	0.31	0.30	1.38	0.01	8.49	0.04	0.13	1.50	6.39	5.69	1.02e <sup>-5</sup>	0.004

\*\*\*, \*\*, \* and ns: significant at 0.1%, 1%, 5% level of probability and non-significant, respectively.

**Table 2.** Phenotypic (CVp) and genotypic (CVg) coefficient of variation and broadcast heritability (H<sub>b</sub><sup>2</sup>) in 30 spring wheat genotypes.

Year	Statistic	Stem No.	Fertile stem No.	Flag leaf length	Flag leaf width	Leaf area	Main stem leaf No.	Spike length	Days to 50% heading	1000 grain weight	Chlorophyll content	Grain iron	Grain zinc
		2009	CVp (%)	18.55	16.66	12.62	9.20	18.48	10.35	8.03	3.32	10.03	7
	CVg (%)	15.41	12.56	10.11	7.96	13.76	8.17	7.49	3.26	9.54	6.01	15.31	9.96
	H <sub>b</sub> <sup>2</sup>	0.69	0.57	0.64	0.75	0.55	0.62	0.87	0.96	0.91	0.74	0.74	0.61
2010	CVp (%)	22.08	22.09	15.41	9.03	16.44	6.29	8.97	4.46	13.17	6.64	14.99	30.72
	CVg (%)	21.67	21.39	15.03	8.67	14.92	5.84	8.78	4.34	12.50	6.03	13.85	29.47
	H <sub>b</sub> <sup>2</sup>	0.96	0.94	0.95	0.92	0.82	0.86	0.96	0.95	0.90	0.83	0.85	0.92

**Table 3.** Descriptive statistics of studied traits in 30 spring wheat genotypes.

Year	Statistic	Stem No.	Fertile stem No.	Flag leaf length	Flag leaf width	Leaf area	Main stem leaf No.	Spike length	Days to 50% heading	1000 grain weight	Chlorophyll content	Grain iron	Grain zinc
		2009	Mean	12.58	6.93	18.41	1.72	18.66	4.32	10.84	77.50	30.89	46.8
	Min	7.20	3.60	11.20	1.34	9.00	2.80	8.40	73.00	22.80	34.96	17.50	22.36
	Max	20.40	11.00	25.80	2.16	30.00	5.80	13.70	81.00	41.6	56.80	48.65	52.69
	CV%	23.8	22.80	16.80	11.20	25.18	13.9	8.90	3.40	10.91	9.30	22.39	16.74
2010	Mean	7.63	5.72	20.13	1.76	24.37	5.16	11.36	67.71	35.07	49.83	30.53	42.45
	Min	4.00	3.20	11.30	1.30	12.50	5.00	8.80	61.00	17.74	37.72	22.74	20.74
	Max	12.00	9.40	27.58	2.16	35.00	6.00	13.90	75.00	48.44	56.84	45.60	78.60
	CV%	22.64	23.70	17.09	11.71	19.01	7.07	9.46	4.67	14.29	7.65	16.87	32.70

#### Estimates of genetic parameters

The phenotypic coefficient of variation ranged from 3.32 to 18.55 in 2009 and from 4.46 to 30.72 in 2010 and the genotypic coefficient of variation ranged from 3.26 to 15.41 in 2009 and from 4.34 to 29.47 in 2010 (Table 2). The highest heritability was exhibited by days to 50% heading in 2009 and by stem number, spike length in 2010 (Table 2).

#### Cluster analysis

Based on cluster analysis, thirty genotypes were grouped into four distinct clusters that group number of genotypes was shown in Fig. 1 suggesting considerable amount of genetic diversity in the material. The accuracy of grouping was confirmed by discriminant analysis and multivariate analysis of variance based on complete randomized design so

that the genotypes correctly belonged in to their groups and the difference between group means were significant ( $P < 0.001$ ) based on Wilks' Lambda, Pillai's trace and Roy's Largest Root tests. The data

presented in Table 4 showed significant ( $P < 0.001$ ) difference among group means for all traits except for main stem leaf number, chlorophyll content, grain iron and grain zinc content.

**Table 4.** Analysis of variance based on Wilks' Lambda statistic.

Statistic	Stem No.	Fertile stem No.	Flag leaf length	Flag leaf width	Leaf area	Main stem No.	Spike length	Days to 50% heading	1000 grain weight	Chlorophyll content	Grain iron	Grain zinc
Wilks' Lambda	0.40***	0.34***	0.52***	0.42***	0.53***	0.93 <sup>ns</sup>	0.46***	0.90 <sup>nd</sup>	0.67**	0.82 <sup>ns</sup>	0.89 <sup>ns</sup>	0.84 <sup>ns</sup>

DF<sub>1</sub>=3, Degree of freedom between groups; DF<sub>2</sub>= 26, Degree of freedom within groups; \*\*\*, \*\* and ns: significant at 0.1%, 1% level of probability and non-significant, respectively.

**Table 5.** Results of group mean comparison and some statistics of clusters.

Cluster	Statistic	Stem No.	Fertile stem No.	Flag leaf length	Flag leaf width	Leaf area	Main stem No.	Spike length	Days to 50% heading	1000 grain weight	Chlorophyll content	Grain iron	Grain zinc
1	mean	10.48 <sup>b</sup>	7.01 <sup>a</sup>	17.98	1.73 <sup>b</sup>	19.86 <sup>b</sup>	4.77 <sup>a</sup>	10.57 <sup>b</sup>	72.92 <sup>a</sup>	24.72 <sup>a</sup>	48.50 <sup>ab</sup>	31.44 <sup>a</sup>	39.44 <sup>a</sup>
	DfTM	0.37	0.69	-1.29	-0.01	-1.66	0.03	-0.53	0.29	0.50	0.18	0.77	0.35
	SD	0.64	0.35	1.63	0.13	1.65	0.19	0.37	2.53	0.59	2.37	2.26	4.39
2	mean	8.93 <sup>c</sup>	5.72 <sup>b</sup>	17.36	1.54 <sup>c</sup>	19.82 <sup>b</sup>	4.81 <sup>a</sup>	10.28 <sup>b</sup>	71.50 <sup>a</sup>	22.55 <sup>b</sup>	48.12 <sup>ab</sup>	29.31 <sup>a</sup>	42.53 <sup>a</sup>
	DfTM	-1.17	-0.61	-1.91	-0.20	-1.70	0.07	-0.82	-1.12	-1.67	-0.20	-1.35	3.45
	SD	0.79	0.48	2.04	0.06	2.59	0.27	0.65	2.82	1.68	2.51	1.73	7.26
3	mean	12.20 <sup>a</sup>	7.43 <sup>a</sup>	20.58 <sup>a</sup>	1.89 <sup>a</sup>	23.57 <sup>a</sup>	4.79 <sup>a</sup>	11.80 <sup>a</sup>	74.33 <sup>a</sup>	25.81 <sup>a</sup>	50.67 <sup>a</sup>	32.78 <sup>a</sup>	41.20 <sup>a</sup>
	DfTM	2.09	1.11	1.30	0.15	2.05	0.05	0.70	1.71	1.60	2.35	2.12	2.11
	SD	0.37	0.21	1.36	0.06	2.13	0.32	0.62	2.92	1.59	2.64	3.64	4.41
4	mean	9.75 <sup>bc</sup>	5.80 <sup>b</sup>	20.65 <sup>a</sup>	1.79 <sup>ab</sup>	22.79 <sup>a</sup>	4.66 <sup>a</sup>	11.63 <sup>a</sup>	72.42 <sup>a</sup>	24.18 <sup>ab</sup>	47.52 <sup>b</sup>	30.11 <sup>a</sup>	36.42 <sup>a</sup>
	DfTM	-0.36	-0.52	1.38	0.06	1.27	-0.07	0.53	-0.21	-0.03	-0.80	-0.55	-2.66
	SD	1.07	0.68	1.53	0.10	1.29	0.25	0.72	2.79	1.69	2.10	4.26	6.57

DfTM: Deviation from total mean; SD: Standard Deviation.

Cluster 1 included 26.66% of genotypes (Fig. 1) and mean of flag leaf length, flag leaf width, leaf area and Spike length were lower than total mean and for other traits were higher than total mean (Table 5). Cluster 1 had the least average distance from centroid and within cluster sum of square for this cluster was 41.81 (Table 6). Cluster 2 included 20% of genotypes and mean of all traits except main stem leaf number and grain zinc were lower than total mean (Table 5). Cluster 2 had 2.70 and 3.40 average and maximum distance from centroid respectively and had 47.44 within cluster sum of square (Table 6). Cluster 3 included 13.33% of genotypes and mean of all traits in this cluster were higher than total mean and had the

highest value for all traits except for grain zinc (Table 5). Cluster 3 had the most average and maximum distance from centroid and within cluster sum of square (Table 6). Also based on mean comparison test, cluster 3 had a rank for all traits. Cluster 4 included 40% of genotypes indicating overall genetic similarity among them and mean of this cluster in more cases were lower than total mean. Cluster 4 had the least maximum distance from centroid and within cluster sum of square (Table 6). The most squared Euclidean genetic distance (75.81) observed between Niknejad and Roshan genotypes. The least squared Euclidean genetic distance (2.71) observed between Shiroodi and Tajan genotypes.

**Table 6.** Between cluster distance, within cluster sum of square and maximum and average distance from centroid for 30 spring wheat genotypes.

	Squared distance between clusters			Within cluster sum of squares	Average distance from centroid	Maximum distance from centroid
	Cluster2	Cluster3	Cluster4			
Cluster1	35.66	43.34	62.34	41.81	2.18	3.07
Cluster2		109.78	59.23	47.44	2.70	3.40
Cluster3			44.92	108.63	2.91	4.12
Cluster4				25.84	2.50	2.88

**Table 7.** between cluster distances and some statistics of clusters for 30 spring wheat genotypes.


	Squared distance between clusters			Within cluster sum of squares	Average distance from centroid	Maximum distance from centroid	Observation No.	Grain iron	Grain zinc
	Cluster2	Cluster3	Cluster4						
Cluster1	7.35	14.45	8.20	26.72	2.15	3.41	5	35.98	41.10
Cluster2		16.75	10.54	97.96	3.12	4.41	9	30.80	46.31
Cluster3			6.51	15.60	1.82	2.89	4	26.61	30.09
Cluster4				133.68	2.89	7.65	12	29.69	35.83

## Discussion

### Analysis of variance and means comparison

Variation among genotypes is appropriate to breeding programs for studied traits. The results presented in Table 1 indicate that. Similarities between results of both years indicate that the trial is replicable and its results are reliable. Results of previous studies on wheat are in accordance with this research, Fareed Khan *et al.* (2004) and Shafeeq *et al.* (2006) for stem number Kashif and Khaliq (2004) for fertile stem number, leaf area and 1000 grain weight, Heidari *et al.* (2006) for flag leaf length, flag leaf width and spike length, days to 50% heading, Morganov *et al.* (2007) for grain iron and zinc concentration. In the previous studies obtained means were 12.72 for fertile stem number, 53.61 for flag leaf area, 14.40 for spike length and 37.96 for 1000 grain weight (Kashif and Khaliq, 2004); 162 for days to 50% heading and 38.3 for 1000 grain weight (Aycicek and Yildirim, 2006), 25.4 for flag leaf length and 2.03 for flag leaf width (Dere and Yildirim, 2006), 46.84 for chlorophyll content (Awaad *et al.*, 2010); 38.0, 31.6 and 37.2 for grain iron and 28, 35.0 and 30.7 for grain zinc content (Welch and Graham 2002; Morgounov *et al.*, 2007) that in more cases, obtained results are in accordance and in a few cases are in contrast with

those. Table 2 content suggesting that environment had no influence on the traits under study especially in 2010.



**Fig. 1.** Dendrogram showing the genetic relationships among 30 spring wheat using ward method and squared Euclidean distance, C1, C2, C3 and C4 contain members of clusters 1, 2, 3 and 4 respectively.

### Heritability

Heritability of all studied traits in both years was high therefore environmental conditions have little impact on the phenotypic differences observed in the studied population (Klug and Cummings, 2005) and these traits can be improved by breeding programs (Courtney, 2007). Heritability of studied traits in this study was estimated by some researchers previously; Kashif and Khaliq (2004) and Ajmal *et al.* (2009)



estimated 0.41 and 0.98 for stem number respectively, Kahrizi *et al.* (2010) estimated 0.43 for stem number and 0.50 for flag leaf length, Smocek (1970) estimated 0.88 for flag leaf length and 0.94 for flag leaf width, Ejaz-ul-hassan and Khaliq (2008) estimated 0.88 for flag leaf area, Dere and Yildirim (2006) estimated 0.5 for flag leaf length and 0.48 for flag leaf width, Memon (2007) estimated 0.93 for fertile stem number and 0.64 for spike length, Ali *et al.* (2008) estimated 0.99 for 1000 grain weight, 0.76 for spike length and 0.46 for fertile stem number, Zhang *et al.* (2009) estimated 0.72 for chlorophyll *a* and 0.75 for chlorophyll *b* content and 0.76 for days to heading time, Awaad (2010) estimated 0.70 for chlorophyll content and 0.76 for flag leaf area and Ghai *et al.* (1969) estimated 0.84 for chlorophyll content. In more cases, obtained results are in accordance with previous studies and in a few cases in contrast with those.

#### *Breeding strategies based on cluster analysis*

Members of the second cluster had the highest grain zinc content. Therefore, genotypes of this cluster can be used in breeding programs with aim of grain zinc improving. As suggested by Kumar *et al.* (2009) who said hybridization between genotypes of divergent cluster will lead to accumulation of suitable genes in a single variety and also suggested to create variability for creating the varieties including a large number of different lines instead of closely related ones, the genotypes of the third cluster may be involve in a multiple crossing program to recover transgressive segregants with high genetic potential of studied traits especially for chlorophyll content, grain iron and grain zinc all together.

Because of the most squared Euclidean genetic distance between Niknejad and Roshan located in cluster 2 and cluster 3 respectively (Fig. 1) and these clusters had the most distance from together. Suggesting by Ilker *et al.* (2010) these genotypes can be used to plan wide crosses, to reach genetic diversity and maximize expression of heterosis. Shiroodi and Tajan genotypes had the least squared Euclidean genetic distance with together and can be

used in back cross program for gene transfer. The frequency distribution of genetic distance value for all 435 pairs of comparisons indicates that 95 % of the pair comparisons had values between 6.3 and 49.4. Only 10 pairs accounting 2.30 % of the total had genetic distance value smaller than 6.3 and 11 pair comparisons, which are only 2.53 % of the total, had values greater than 49.4. Cophenetic value of 0.75 or more are usually recommended for the best fit of the cluster analysis (Ribeiro Trindade *et al.*, 2010). The present study confirmed a good fit with the genetic distance matrix by having cophenetic correlation coefficient of  $r=0.793$ .

Because of grain iron and grain zinc importance and presence meaningful genetic diversity among genotypes for these traits (Table 1), genotypes were grouped based on these traits. The most squared Euclidean genetic distance (556.45) observed between Mahdavi and Bahar. Therefore, these genotypes can be used as parents in hybridization program. The least squared Euclidean genetic distance (0.07) observed between Shiroodi and Tajan genotypes. Some information about clusters, which extracted through cluster analysis based on grain iron and zinc content, presented in Table 7.

#### **References**

**Ajmal SU, Zakir N, Mujahid MY.** 2009. Estimation of genetic parameters and character association in wheat. *Journal of Agricultural and Biological Science* **1(1)**, 15-18.

**Akhtar S, Ashgar A.** 2011. Mineral fortification of whole wheat flour: An overview. In: flour and breads and their fortification in health and disease prevention (Preedy, V.R., Watson, R.R. and Patel, V.B. eds), San Diego: Academic Press. p. 263-271.

<http://dx.doi.org/10.1016/B978-0-12-380886-8.10024-8>

**Ali Y, Manzoor-Atta B, Akhter J, Monneveux P, Lateef Z.** 2008. Genetic variability, association and diversity studies in wheat (*Triticum aestivum* L.) Germplasm. *Pakistan Journal of Botany* **40(5)**, 2087-2097.



- Awaad HA, Youssef MAH, Moustafa ESA.** 2010. Identification of genetic variation among bread wheat genotypes for lead tolerance using morpho – physiological and molecular markers. *Journal of American Science* **6(10)**, 1142-1153.
- Bouis HE, Graham RD, Welch RM.** 2000. The Consultative Group on International Agriculture Research (CGIAR) Micronutrients Project: justification and objectives. *Food and Nutrition Bulletin* **21(4)**, 374–381.
- Burton GW.** 1952. Quantitative inheritance in grasses. In: *Proceeding of the 6<sup>th</sup> International Grassland Congress*, vol. 1, Washington DC, pp. 277-283.
- Cakmak I, Torun A, Millet E, Feldman M, Fahima T, Korol A, Nevo E, Braun HJ, Özkan H.** 2004. *Triticum dicoccoides*: An important genetic resource for increasing zinc and iron concentration in modern cultivated wheat. *Soil Science and Plant Nutrition* **50(7)**, 1047-1054.  
<http://dx.doi.org/10.1080/00380768.2004.10408573>
- Cakmak I.** 2008. Enrichment of cereal grains with Zn: Agronomic or genetic biofortification. *Plant and Soil* **302**,1–17.  
<http://dx.doi.org/10.1007/s11104-007-9466-3>
- Courtney MW.** 2007. Genotypic variability and inheritance of iron and zinc in sweet potato. MS Thesis. Agricultural and Mechanical College, Louisiana State University.
- Dere S, Yildirim MB.** 2006. Inheritance of grain yield per plant, flag leaf width, and length in an 8 × 8 diallel cross population of bread wheat (*T. Aestivum* L.). *Turkish Journal of Agriculture and Forestry* **30**, 339-345.
- Ejaz-U-Hassan S, Khaliq I.** 2008. Quantitative inheritance of some physiological traits for spring wheat under two different population densities. *Pakistan Journal of Botany* **40(2)**, 581-587.
- Fareed Khan M, Khalid Khan M, Kazmi M.** 2004. Genetic variability among wheat cultivars for yield and yield components under the agro-ecological conditions of district rawalakot, Azad Kashmir, Pakistan. *Sarhad Journal of Agriculture* **20**, 391-395.
- Ghai BS, Srivastava AK, Gill KS.** 1969. Inheritance of amount of chlorophyll in wheat, *triticum aestivum* L. *Euphytica* **18**, 403-405.  
<http://dx.doi.org/10.1007/BF00397789>
- Gupta PK.** 2000. Soil, plant, water and fertilizer analysis. Agrobios, New Dehli, India.
- Heidari B, Saiddi GA, Badredin EST, Sounaga K.** 2006. Study of genetic variation and estimate heritability of some quantitative traits in double haploid wheat lines. *Iranian Journal of Agricultural Sciences* **37(2)**, 347-356.
- Ilker E, Aykut Tonk F, Tosun M.** 2010. Heterosis for yield and its components in bread wheat crosses among powdery mildew resistant and susceptible genotypes. *Pakistan Journal of Botany* **42(1)**, 513-522.
- Johnson HW, Robinson HF, Comstock RE.** 1955. Estimates of genetic and Environmental variability in Soybeans. *Agronomy Journal* **47(7)**, 314-318.  
<http://dx.doi.org/10.2134/agronj1955.00021962004700070009x>
- Joshi BK, Mudwari A, Bhatta MR, Ferrara GO.** 2004. Genetic diversity in Nepalese wheat cultivars based on agro-morphological traits and coefficients of parentage. *Nepal Agriculture Research Journal* **5**, 7-17.
- Kahrizi D, Cheghamirza K, Kakaei M, Mohammadi R, Ebadi A.** 2010. Heritability and genetic gain of some morphophysiological variables of

durum wheat (*Triticum turgidum* var. durum). African Journal of Biotechnology **9**(30), 4687-4691.

**Kashif M, Khaliq I.** 2004. Heritability, correlation and path coefficient analysis for some metric traits in wheat. International Journal of Agriculture and Biology **6**, 138-142.

**Klug WS, Cummings MR.** 2005. Essentials of genetics. 5<sup>th</sup> ed. Upper Saddle River, New Jersey. Pearson Education, Inc.

**Kumar B, Lal GM, Ruchi Upadhyay A.** 2009. Genetic variability, diversity and association of quantitative traits with grain yield in bread wheat (*Triticum Aestivum* L.). Asian Journal of Agricultural Sciences **1**(1), 4-6.

**Memon S, Qureshi MUD, Ansari BA, Sial MA.** 2007. Genetic heritability for grain yield and its related characters in spring wheat (*Triticum aestivum* L.). Pakistan Journal of Botany **39**(5), 1503-1509.

**Mishra RS, Lotha RE, Mishra SN, Paul PK, Mishra HN.** 1988. Results of heterosis breeding on chilli (*Capsicum annum* L.). Capsicum Newsletter **7**, 49-50.

**Mohammadi SA, prasanna BM.** 2003. Analysis of genetic diversity in crop plants: salient statical tools and considerations. Crop Science **43**, 1235-1248.

<http://dx.doi.org/10.2135/cropsci2003.1235>

**Morgounov A, Gomez-Becerra HF, Abugalieva A, Dzhunusova M, Yessimbekova M, Muminjanov H, Zelenskiy Y, Ozturk L, Cakmak I.** 2007. Iron and zinc grain density in common wheat grown in Central Asia. Euphytica **155**, 193-203.

<http://dx.doi.org/10.1007/s10681-006-9321-2>

**Ranum P.** 2001. Zinc enrichment of cereal staples. Food and Nutrition Bulletin **22**, 169-172.

**Ribeiro Trindade AP, Barth Pinto RJ, Amaral Júnior AT, Mangolin AC, Silva Machado MFP, Scapim CA.** 2010. Genetic diversity of breeding popcorn lines determined by SSR markers. Electronic Journal of Biotechnology **13**(1), 1-9.

<http://dx.doi.org/10.2225/vol13-issue1-fulltext-11>

**Sayar R, Khemira H, Kharrat M.** 2007. Inheritance of deeper root length and grain yield in half-diallel durum wheat (*Triticum durum* Desf.) crosses. Annals of Applied Biology **151**, 213-220.

<http://dx.doi.org/10.1111/j.1744-7348.2007.00168.x>

**Shafeeq S, Rahman MU, Zafar Y.** 2006. Genetic variability of different wheat (*Triticum aestivum* L.) Genotypes/cultivars under induced water stress. Pakistan Journal of Botany **38**(5), 1671-1678.

**Singh RK, Chowdhury BD.** 1985. Biometrical method in quantitative genetic analysis. Kalyani publishers, Ludhiana, New Delhi, 54-57 p.

**Small H, Stevens TS, Bauman WC.** 1975. Novel ion exchange chromatographic method using conductimetric detection. Analytical Chemistry **47**, 1801-1809.

**Smocek J.** 1970. Prediction of relative efficiency of some selection indices used in winter wheat. Biologia Plantarum **12**(3), 216-223.

<http://dx.doi.org/10.1007/BF02920870>

**Toker C, Cagirgan MI.** 2004. The use of phenotypic correlation and factor analysis in determining characters for grain yield selection in chickpea (*Cicer arietinum* L.). Hereditas **140**, 226-8.

<http://dx.doi.org/10.1111/j.1601-5223.2004.01781.x>

**Welch RM, Graham RD.** 2002. Breeding crops for enhanced micronutrient content. Plant and Soil **245**, 205-214.

<http://dx.doi.org/10.1023/A:1020668100330>

**Welch RM, Graham RD.** 2004. Breeding for micronutrients in staple food crops from a human

nutrition perspective. *Journal of Experimental Botany* **55**, 353-364.

<http://dx.doi.org/10.1093/jxb/erh064>

**White PJ, Broadley MR.** 2005. Biofortifying crops with essential mineral elements. *Trends in Plant Science* **10**, 586–593.

<http://dx.doi.org/10.1016/j.tplants.2005.10.001>

**Zhang K, Zhang Y, Chen G, Tian J.** 2009. Genetic analysis of grain yield and leaf chlorophyll content in common wheat. *Cereal Research Communications* **37(4)**, 499–511.

<http://dx.doi.org/10.1556/CRC.37.2009.4.3>