



## RESEARCH PAPER

## OPEN ACCESS

## Evaluation of essential oil production of dill affected by different intercropping patterns with vetch and harvesting times

Parviz Karimzadeh<sup>1\*</sup>, Saeid Zehtab-Salmasi<sup>1</sup>, Jalil Shafagh-Kalvanagh<sup>1</sup>, Hossein Janmohammadi<sup>2</sup>

<sup>1</sup>Department of Plant Eco-Physiology, Faculty of Agriculture, University of Tabriz, Iran

<sup>2</sup>Department of Animal Science, Faculty of Agriculture, University of Tabriz, Iran

**Key words:** Additive series, dill, essential oil, harvesting times, intercropping, vetch.

<http://dx.doi.org/10.12692/ijb/4.9.20-25>

Article published on May 07, 2014

### Abstract

Intercropping is considered as the practical application of ecological principles such as diversity, interference and other natural regulation mechanisms. In this research vetch and dill were intercropped at different additive series (100:25, 100:50 and 100:75) at the Research Farm of the Faculty of Agriculture, University of Tabriz, Iran. Field experiment was arranged as split plot based on randomized complete block design with three replicates. Dill umbels were harvested at flowering, dough development and complete ripening stages. Results showed that among harvesting times, dough development stage and among intercropping patterns, 100:25 treatment had the highest essential oil percentage, essential oil yield and harvest index of essential oil. Vetch as a forage and legume crop promotes dill essential oil production and could be an effective plant in intercropping systems with essential oil bearing medicinal plants such as dill.

\* **Corresponding Author:** Parviz Karimzadeh ✉ [p.k\\_parviz@yahoo.com](mailto:p.k_parviz@yahoo.com)

## Introduction

Dill (*Anethum graveolens* L.) is an annual and sometimes biennial herb, which is native south-west Asia and south-east Europe and has been cultivated since ancient times (Bailer *et al.*, 2001). The major components of *A. graveolens* are flavonoids, phenolic compounds and essential oil (Amin and Sleem, 2007). Many studies were interested in the chemical composition of essential oil of *A. graveolens*, such as its antibacterial (Delaquis *et al.*, 2002; Jirovetz *et al.*, 2003; Raffi and Shahverdi, 2006), antifungal activity (Tian *et al.*, 2012), insecticidal properties (Chaubey, 2007; Seo *et al.*, 2009) and its essence has an inhibitory effect on stored potatoes sprouting (Zehtab-Salmasi *et al.*, 2006). Carrubba *et al.*, (2008) indicated that the presence of dill exerted residue in the soil had a significant effect on fennel seed yield at following years.

Vetch has a considerable potential to cultivate as cover or forage crop especially in some west Asian regions (Abd El Moneim and Elias, 2003). This crop also has nitrogen biological fixation ability and can be effective to soil fertility and reduction of fertilizer N input (Dabney *et al.*, 2001). Therefore, in pure or mixture cultivation it can provide the rotational benefits (Twidwell *et al.*, 1987).

Plant mixtures can be formed by adding together the plant populations used in the pure stands (Agboola and Fayemi, 1971). This means that in such additive intercropping systems the total plant population of the mixtures is doubled when two crops are intercropped in this manner (Ebwongu *et al.*, 2001). In other words, an inherent feature of additive intercropping is that the total plant population of the mixture is greater than that of the pure stands, which may contribute to its yield advantage (Willey and Osiru, 1972).

The biosynthesis of secondary metabolites, although controlled genetically, it is strongly affected by environmental factors, and also by agronomic conditions, harvesting time and the type of processing (Miguel *et al.*, 2004). One of the most important

characteristics of oil essential accumulation is its dependence on the developmental stage of the plant per se as well as its concerned part (Sangwan *et al.*, 2001).

Some further interest in the potential role of medicinal and aromatic plants in intercropping systems has arisen from the widespread trend toward the cultivation of these plants. Nitrogen fixation by some plants may improve essential oil yield of medicinal plants in intercropping. Thus, this research was to evaluate the effect of different intercropping patterns and different harvesting times on essential oil of dill intercropped with vetch.

## Materials and methods

The field experiment was conducted at the Research Farm of the Faculty of Agriculture, University of Tabriz, Iran (38°5N, 46°E) at 2013. The experiment was arranged as split plot on the basis of randomized complete block design with three replicates. In this study dill (*Anethum graveolens* L.) and vetch (*Vicia villosa*) intercropped at different additive series (100:25, 100:50 and 100:75). The main and secondary crops were vetch and dill, respectively. As dill seeds are sensitive to seed bed conditions, therefore soft and smooth seed bed prepared and covered by thin layer of sand silt. Each plot consisted of 10 rows and seeds were sown at about 1-3 cm deep. Optimum density for dill and vetch were considered as 100 and 200 plant/m<sup>2</sup>, respectively.

### Field conditions and treatments

Dill umbels were harvested at flowering, dough development and complete ripening stages. At each stage, 10 plants were harvested from each plot and the umbels were separated and dried. Then, dry umbels were distilled with 500 ml water and hydro-distillation was continued for 3 hours, using a Clevenger-type apparatus.

### Data collection

All data were analyzed using MSTAT-C statistical package (MSTAT, 1993). Means of each trait were compared according to Duncan multiple range test at

$p \leq 0.05$ .

## Results and discussion

### Statistical analysis

Analysis of variance of the data showed that the effect of different intercropping patterns (main factor), harvesting times (secondary factor), and their interaction were significant on essential oil percentage and essential oil yield and harvest index of

essential oil (Table 1). Almost in all of treatments intercropped dill plants produced a significantly higher amount of essential oil when compared to monocultured plants (Table 2). It seems that nitrogen fixation by vetch helped dill plants to improve essence production. Similarly, Maffei and Mucciarelli, (2003) relived that in peppermint/soybean strip intercropping, essential oil content increased up to 50% and quality of essential oil improved.

**Table 1.** Analysis of variance of the data for essential oil percentage, yield and harvest index of dill at different harvesting times and intercropping patterns with vetch.

Source	df	Characters		
		Essential oil percentage	Essential oil yield	Essential oil harvest index
Replication	2	0.028	0.107	0.001
Intercropping patterns (I)	3	0.873**	6.655**	0.294**
Error (I)	6	0.039	0.111	0.008
Harvest time (T)	2	4.73**	46.472**	3.915**
I×T	6	0.18**	2.482**	0.12**
Error (T)	16	0.037	0.087	0.005
Total	35			
C.V%		13.40	14.26	11.04

\*\* Significant at  $p \leq 0.01$ .

There were no significant differences among intercropping patterns at flowering stage, and 100:50 treatment had the highest essential oil percentage (0.991%). The highest essential oil yield and harvest index of essential oil were obtained from 100:75 and sole dill treatments, respectively. At dough development stage, 100:25 treatment had the highest

essential oil percentage (2.838%). Also in this stage the highest yield and harvest index of essential oil were obtained from 100:25 treatment. At complete ripening stage, the highest and lowest essential oil percentage, essential oil yield and harvest index of essential oil were recorded for 100:25 and sole dill treatments, respectively (Table 2).

**Table 2.** Mean of the essential oil percentage, essential oil yield and essential oil harvest index of dill for harvesting times and intercropping patterns with vetch.

Harvesting time	Treatment	Characters		
		Essential oil percentage (%)	Essential oil yield (g/m <sup>2</sup> )	Essential oil harvest index (%)
Flowering stage	100:25	0.947 de	0.385 f	0.115 g
	100:50	0.991 de	0.348 f	0.109 g
	100:75	0.945 de	0.39 f	0.116 g
	Sole dill	0.746 e	0.362 f	0.135 g
dough development stage	100:25	2.838 a	6.777 a	1.8 a
	100:50	2.104 b	4.232 b	1.188 b
	100:75	2.033 b	3.142 c	0.943 c
	Sole dill	1.562 c	2.764 c	1.021 c
Full ripening	100:25	1.796 bc	2.706 c	0.711 d
	100:50	1.219 d	1.536 d	0.455 e
	100:75	1.12 de	1.369 de	0.422 ef
	Sole dill	1.026 de	0.879 ef	0.325 f

Different letters in each column indicate significant difference at  $p \leq 0.05$ .

Essential oil production at flowering stage was lower than that at dough development and maturity stages (Table 2). In contrast, İNAN *et al.*, (2011) showed that maximum essential oil percentage (3.56 %) for *Thymbra spicata* were obtained full flowering stage. Yoshida and Sawasaki, (1978) found that the maximum clary sage oil content was obtained at the end of the blossom period. In other research Mirza *et al.*, (2011) indicated that the highest percentage of essential oil (2.8%) of *Mentha piperita* was recorded in full flowering stage, while *Salvia officinalis* had the highest percentage of essential oil (3%) in early flowering stage.

The highest essential oil percentage, yield and harvest index were recorded for dough development stage, followed by maturity stage (Table 2). Similarly, Callan *et al.*, (2007) reported the highest oil yields of dill were obtained when most of the fruits on primary umbels were pigmented but had not become dry and fully mature. Aidi Wannes *et al.*, (2009) indicated that the highest fruit essential oil of *Myrtus*

**Abd El Moneim AM, Elias SF.** 2003. Underground vetch (*Vicia sativa* ssp. *amphicarpa*): A potential pasture and forage legume for dry areas in West Asia. *Journal of Agronomy and Crop Science* **189**, 136–141.

**Agboola AA, Fayemi AA.** 1971. Preliminary trials on the intercropping of maize with different tropical legumes in Western Nigeria. *The Journal of Agricultural Science* **77**, 219–225.

<http://dx.doi.org/10.1017/S0021859600024345>

**Aidi Wannes W, Mhamdi B, Marzouk B.** 2009. Variations in essential oil and fatty acid composition during *Myrtus communis* var. *italica* fruit maturation. *Food chemistry* **112**, 621–626.

<http://dx.doi.org/10.1016/j.foodchem.2008.06.018>

**Amin WM, Sleem AA.** 2007. Chemical and biological study of aerial parts of dill (*Anethum*

*communis* was obtained 60 days after flowering stage. Telci *et al.*, (2006) showed that 20 days after flowering of *Coriandrum sativum* maximum essential oil content achieved. Arif *et al.*, (2010) in sea buckthorn berries study reported that essential oil content increased by fruit maturity.

Finally, Present study suggests that harvesting time (flowering, dough development and complete ripening stages) and intercropping patterns had significant effect on essential oil production of dill; therefore these two factors important in essential oil production of Aromatic Plants. In this research the best harvesting time and intercropping patterns were dough development stage and 100:25 treatment, respectively. Also it could be suggested that intercropping of dill with nitrogen fixing plants such as vetch can enhance essential oil production per unit area.

## References

*graveolens* L.). *Egyptian Journal of Biomedical Sciences* **23**, 73–90.

<http://dx.doi.org/10.4314/ejbs.v23i1.40296>

**Arif S, Ahmed SD, Shah AH, Hassan L, Awan SI, Hamid A, Batool F.** 2010. Determination of optimum harvesting time for vitamin C, oil and mineral elements in berries sea buckthorn (*Hippophae rhamnoides*). *Pak J Bot* **42**, 3561–3568.

**Bailer J, Aichinger T, Hackl G, de Hueber K, Dachler M.** 2001. Essential oil content and composition in commercially available dill cultivars in comparison to caraway. *Industrial crops and products* **14**, 229–239.

[http://dx.doi.org/10.1016/S0926-6690\(01\)00088-7](http://dx.doi.org/10.1016/S0926-6690(01)00088-7)

**Callan NW, Johnson DL, Westcott MP, Welty LE.** 2007. Herb and oil composition of dill (*Anethum graveolens* L.): Effects of crop maturity and plant density. *Industrial Crops and products* **25**, 282–287.

<http://dx.doi.org/10.1016/j.indcrop.2006.12.007>

- Carrubba A, la Torre R, Saiano F, Aiello P.** 2008. Sustainable production of fennel and dill by intercropping. *Agronomy for sustainable development* **28**, 247–256.
- Chaubey MK.** 2007. Insecticidal activity of *Trachyspermum ammi* (Umbelliferae), *Anethum graveolens* (Umbelliferae) and *Nigella sativa* (Ranunculaceae) essential oils against stored-product beetle *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae). *African Journal of Agricultural Research* **2**, 596–600.
- Dabney SM, Delgado JA, Reeves DW.** 2001. Using winter cover crops to improve soil and water quality. *Communications in Soil Science and Plant Analysis* **32**, 1221–1250.  
<http://dx.doi.org/10.1081/CSS-100104110>
- Delaquis PJ, Stanich K, Girard B, Mazza G.** 2002. Antimicrobial activity of individual and mixed fractions of dill, cilantro, coriander and eucalyptus essential oils. *International journal of food microbiology* **74**, 101–109.  
[http://dx.doi.org/10.1016/S0168-1605\(01\)00734-6](http://dx.doi.org/10.1016/S0168-1605(01)00734-6)
- Ebwongu M, Adipala E, Ssekabembe CK, Kyamanywa S, Bhagsari AS.** 2001. Effect of intercropping maize and solanum potato on yield of the component crops in central Uganda. *African Crop Science Journal* **9**, 83–96.
- İNAN M, KIRPIK M, KAYA DA, KIRICI S.** 2011. Effect of harvest time on essential oil composition of *Thymbra spicata* L. growing in flora of Adiyaman. *Advances in Environmental Biology* **5**, 356–358.
- Jirovetz L, Buchbauer G, Stoyanova AS, Georgiev E V, Damianova ST.** 2003. Composition, quality control, and antimicrobial activity of the essential oil of long-time stored dill (*Anethum graveolens* L.) seeds from Bulgaria. *Journal of agricultural and food chemistry* **51**, 3854–3857.  
<http://dx.doi.org/10.1021/jf030004y>
- Maffei M, Mucciarelli M.** 2003. Essential oil yield in peppermint/soybean strip intercropping. *Field crops research* **84**, 229–240.  
[http://dx.doi.org/10.1016/S0378-4290\(03\)00092-3](http://dx.doi.org/10.1016/S0378-4290(03)00092-3)
- Miguel MG, Guerrero C, Rodrigues H, Brito JC, Duarte F, Venâncio F, Tavares R.** 2004. Main Components of the Essential Oils from Wild Portuguese *Thymus mastichina* (L.) L. ssp. *mastichina* in Different Developmental Stages or Under Culture Conditions. *Journal of Essential Oil Research* **16**, 111–114.  
<http://dx.doi.org/10.1080/10412905.2004.9698665>
- Mirza M, Ghoraishi F, Bahadori A.** 2011. Effect of harvesting time on essential oils content and composition of *Salvia officinalis* L. and *Mentha piperita* L. in Khuzestan province. *Iranian j. Medicinal and Aromatic Plants*. **26**, 531–543.
- MSTAT C.** 1993. MSTAT-C, a microcomputer program for design, arrangement and analysis of agronomic research experiments. Michigan State University.
- Raffi F, Shahverdi AR.** 2006. Comparison of essential oils from three plants for enhancement of antimicrobial activity of nitrofurantoin against enterobacteria. *Chemotherapy* **53**, 21–25.  
<http://dx.doi.org/10.1159/000098246>
- Sangwan NS, Farooqi AHA, Shabih F, Sangwan RS.** 2001. Regulation of essential oil production in plants. *Plant Growth Regulation* **34**, 3–21.  
<http://dx.doi.org/10.1023/A:1013386921596>
- Seo S-M, Kim J, Lee S-G, Shin C-H, Shin S-C, Park I-K.** 2009. Fumigant antitermitic activity of plant essential oils and components from ajowan (*Trachyspermum ammi*), allspice (*Pimenta dioica*), caraway (*Carum carvi*), dill (*Anethum graveolens*), geranium (*Pelargonium graveolens*), and litsea

(*Litsea cubeba*) oils against. *Journal of agricultural and food chemistry* **57**, 6596–6602.

<http://dx.doi.org/10.1021/jf9015416>

**Telci I, Bayram E, Avci B.** 2006. Changes in yields, essential oil and linalool contents of *Coriandrum sativum* varieties (var. *vulgare* Alef. and var. *microcarpum* DC.) harvested at different development stages. *European Journal of Horticultural Science* **71**, 267.

**Tian J, Ban X, Zeng H, He J, Chen Y, Wang Y.** 2012. The mechanism of antifungal action of essential oil from dill (*Anethum graveolens* L.) on *Aspergillus flavus*. *PLoS one* **7**, e30147.

<http://dx.doi.org/10.1371/journal.pone.0030147>

**Twidwell EK, Johnson KD, Cherney JH.** 1987. Forage potential of soft red winter wheat-hairy vetch mixtures. *Applied agricultural research (USA)* **2**, 164-169.

**Willey RW, Osiru DSO.** 1972. Studies on mixtures of maize and beans (*Phaseolus vulgaris*) with particular reference to plant population. *The Journal of Agricultural Science* **79**, 517–529.

<http://dx.doi.org/10.1017/S0021859600025909>

**Yoshida T, Sawasaki T.** 1978. On the variations of the percentage yield and the chemical composition of essential oil in clary sage (*Salvia sclarea* L.). *Japanese journal of tropical agriculture* **21**, 145-149.

**Zehtab-Salmasi S, Ghassemi-Golezani K, Moghbeli S.** 2006. Effect of Sowing Date and Limited Irrigation on the Seed Yield and Quality of Dill (*Anethum graveolens* L.). *Turkish Journal of Agriculture & Forestry* **30**, 281-286.