



RESEARCH PAPER

OPEN ACCESS

## Seasonal dynamics of the phytoplankton in relation to environmental variables in Nguru Lake, Northeastern Nigeria

M. M. Abubakar<sup>1\*</sup>, J. Yaji Abubakar<sup>2</sup>

<sup>1</sup>*Department of Biological Sciences, Federal University Dutse, Nigeria*

<sup>2</sup>*Department of Fisheries and Aquaculture, Adamawa State University, Mubi, Nigeria*

Article published on January 21, 2013

**Key words:** Phytoplankton, nutrients, water quality, Nguru lake.

### Abstract

Studies were conducted in Nguru Lake between May 2006 and April 2008 to map temporal changes in phytoplankton composition and attempt to relate these changes to environmental parameters in this ecosystem. Twenty five phytoplankton genera were recorded. Green algae (Chlorophyta) were the most dominant contributing 46% of the total biomass followed by blue green algae-Cyanophyta(29%), diatoms-Bacillariophyta(22%) and finally the Dinophyta(2%). Seasonal changes were observed for the three major plant nutrients total phosphorus (PO<sub>4</sub>-P) ranged from 5.2 -9.6mg/l. Total nitrogen (NO<sub>3</sub>-N) ranged from 2.4-12.8mg/l and sulphates ranged from 1.0-10.1mg/l. wide variations in environmental variables were observed in the lake. Results obtained in this study showed that the lake was undergoing gradual eutrophication resulting in deterioration of water quality, The eutrophication was further aggravated by increasing human and livestock population.

\*Corresponding Author: M. M. Abubakar ✉ [mmabubakar2005@yahoo.com](mailto:mmabubakar2005@yahoo.com)

## Introduction

Water is an important basic resource for humanity. The availability and quality of this resource is however rapidly declining with increase in population.

The quality of water, especially of eutrophic waters is essentially determined by the quantity of algae. Drinking water supply, recreational activities and fisheries can be impaired by high phytoplankton biomass. Therefore, water quality management has to monitor the phytoplankton content in order to come up with preventive measures such as aeration to prevent fish kills during decay of the planktonic biomass (Imhoff and Alberrecht 1975).

Phytoplankton development in aquatic ecosystems is greatly enhanced by increased level of the major plant nutrients. This is because some of the nutrients are involved in the intercellular metabolic regulation and as building blocks in protein molecules (Crul 1993). The major nutrients that regulate algal growth are phosphorus, nitrogen and dissolved silica. Water temperature and turbidity are equally important for they have direct bearing on the development and response of phytoplankton.

Previous documented studies on Nguru Lake include the works of Abdullahi(1997), Lafiagi(1997), WDI(2006) and Abubakar (2007) revealed that the phytoplankton of the lake was dominated by the green algae and to a less extent the blue green algae. In the recent past the lake has been besieged by a myriad of ecological disruptions including among others dumping of fertilizers as a result of agricultural activities thereby leading to frequent algal blooms, excessive siltation, proliferation of aquatic weeds, and draught.

The aim of the present study is to document the temporal changes of the phytoplankton and relate same to the environmental variables.

## Materials and methods

### *Study area*

Nguru Lake is a part of the Hadejia Nguru wetlands (HNW), which are located in northeastern Nigeria. Nguru lake occupies an area of approximately 58.100ha and is located between latitude 12°40'N and 13°60'N and longitudes 10°20'E and 11°00'E. The lake developed from two main drainage systems. The Hadejia and Jama'are rivers. The lake is surrounded by a flood plain made up of a network of channels and pools producing a complete pattern of permanently and seasonally flooded land and dry land (Hollis *et al*, 1993).

The inter-tropical convergence zone (ICZT) influences the climate of the Nguru area. This climate is characterized by distinct wet and dry seasons, which are controlled by the cyclical migration of the ICTZ. Rainfall in this area usually starts from May and end in September or October, in most cases with a long break of up to a month between the first rain and subsequent ones. There is also a wide variation of rainfall even within the wetlands area, from the south where the rains start earlier to the North-east where they occur later, the difference in amount of rainfall ranging from 500mm to about 700mm per annum. Latest records of rainfall show that it has been below average for almost twenty years (Adams, 1993).

Over half a million people depend upon the lake and the surrounding wetlands for their livelihoods, especially for water supply. Lafiagi (1997) has identified about 65 species of fishes. Therefore, majority of the people living around the lake are either fishermen, or processors and marketers of fish. The land around the lake is irrigated in both dry and wet season for vegetables and rice. The lake is also used for livestock grazing, for recreation and research, since it has a large biodiversity of birds, insects, wildlife and flowering plants. In fact the lake and the surrounding wetlands is a particularly good representative of a natural or near-natural wetlands, which embodies all the diverse flora and fauna at

both the Sahel and the Sudan savanna in a single location; hence the area is important for ecotourism. Nguru Lake is designated as a Ramsar site by the Ramsar convention; it is presently the only Ramsar site in Nigeria.

*Sampling for water and phytoplankton*

Samples for water and phytoplankton were taken on monthly basis from May 2006 to April 2008 from five stations. The samples were collected between 07.00am and 11.00am.

Water samples were collected with a Van Dorn sampler. A portion of the water samples were analyzed for total alkalinity by titration with methyl orange and phenolphthalein indicators. Total phosphorus, nitrogen, sulphate and turbidity were determined spectrophotometrically as described by APHA (1985).

Transparency was determined in the field with a 20cm diameter black and white *Secchi* disc.

Temperature, pH and conductivity were measured with pH meter model 3150.

For phytoplankton analysis, 250ml of the water was fixed in the field with Lugol's solution. The sample was then sedimented for 48hours after which enumeration of the phytoplankton was done using Sedgwick-rafter counting chamber, according to the methods of Boyd (1981). Identification of the phytoplankton was done following Prescott (1951).

*Data Analysis*

The seasonal data was analyzed by one-way analysis of variance (ANOVA), and Duncan's Multiple Range Test was used to compare the means of the data collected, Pearson's product moment correlation coefficient was used to determine association between the physico-chemical properties and the biological properties. While similarity between the two years concerning the measured environmental and biotic variables was investigated by the simple Student's t-test. Biological diversity was measured with the Shannon-Weiner index and Evenness index.

**Table 1.** Mean results for physico-chemical properties of Nguru Lake for the two seasons.

Parameter	Means for dry season	Means for rainy season
Temperature	16.83b	22.90a
pH	8.11b	8.65a
Transparency	63.59a	59.33a
Depth	104.36a	105.06a
TDS	80.80b	86.30a
Suspended solids	3.51b	4.52a
Turbidity	6.19a	6.92a
Conductivity	177.69a	174.23b
Alkalinity	38.83a	33.88b
DO	6.89a	6.55b
BOD	4.61a	4.67a
COD	8.65b	10.26a
Total Phosphate	6.77a	6.98a
Total Nitrogen	6.09a	5.57a
Sulphate	3.74b	6.75a
Magnesium	0.07a	0.09a
Calcium	0.17a	0.19a
Iron	0.22a	0.24a
Zinc	0.97a	0.42a
Copper	0.07a	0.07a
Manganese	4.49a	4.14a

Means with the same letter in the same column are not significantly different using DMRT.

**Table 2.** Mean distribution of Phytoplankton in Nguru Lake and the diversity indices of the stations.

Chlorophyta	Station 1	Station 2	Station 3	Station 4	Station 5
Ankistrodesmus falcatus Kom.	****	****	**	**	*
Closterium aciculare West	***	****	**	-	-
Closterium parvulum Nag.	**	***	**	-	-
Closterium sp.	***	-	-	-	-
Spirogyra cormunis Her.	-	-	**	*	*
Pediastrum boryanum Ralfs.	**	*	-	-	-
Pediastrum simplex Ralfs	-	**	-	-	-
Staurastrum leptocladum Kutz	**	**	-	-	-
Zygnema sterile Ehr	**	***	***	*	*
Chlorella vulgaris Breb	**	**	*	*	*
Cyanophyta					
Anabaena circinalis Nag.	*	*	*	****	*
Anabaena flos-aque Nag.	*	*	**	****	**
Anacystis cyanae Bom	-	-	*	**	*
Gomphophaeria sp.	**	***	-	-	-
Microcystis aeruginosa Kirchn.	*	**	*	*	*
Spirulina massarti Nag	-	-	**	**	*
Oscillatoria limnosa Gomont	*	**	*	*	*
Bacillariophyta					
Asterionella Formosa Grun	-	-	-	*	*
Cyclotella bodanica Kutz	-	-	-	*	-
Diatom vulgare Breb	***	-	-	-	-
Navicula radiosa Kutz	*	*	*	-	-
Synedra ulna Mer.	**	**	*	-	-
Nitzschia palea Kutz	**	**	-	*	-
Dinophyta					
Ceratium hirundinella Kom	*	*	-	*	-
Mean No of individuals(N)	11963	12154	10402	13244	2434
Number of species(S)	18	16	14	13	11
Shannon-Weaver index(D)	5.32	3.89	4.67	2.92	2.11
Evenness index(E)	0.9821	0.8762	0.9211	0.6322	0.5412

\*\*\*\*Present in >50% of the samples; \*\*\* Present in > 30% of the samples; \*\*Present in >20% of the samples; \*Present in <10% of the samples

**Results**

The temperature ranged from 7°C to 32°C. In the rainy season (May-Sept) of the first year and in the second year temperature range is 15°C to 32°C, and 17°C to 30°C respectively. While in the dry season of the first and second year, the temperature range is from 9°C to 21°C and 7°C to 22°C respectively. The temperature at Nguru Lake shows highly significant seasonal variation (p < 0.001), there was also highly significant spatial variation (p < 0.001). pH showed a range of 7.6-9.5 with no significant seasonal or spatial variation. Transparency ranged from 11-147cm. The mean turbidity during the study period is 6.49 ± 0.02ml<sup>-1</sup>. Conductivity and alkalinity showed significant seasonal and spatial variation during the study period, both showing maxima in the dry season (Table 1). The nutrients, phosphorus, nitrogen and sulphate exhibited similar pattern, with all showing maxima during the rainy season

mainly due to surface run-off from agricultural lands. All the nutrients showed significant seasonal variation.

Of all the environmental variables examined in this study only the nutrients showed significant correlation with the phytoplankton (r > 0.5).

The phytoplankton of Nguru Lake consists of four genera, with twenty-five species (Table 2). The phytoplankton was dominated by the Chlorophyta, which makes up 46% of the total algae. The Cyanophyta constituted 29% and the Bacillariophyta contributes 22% of the population, while the Dinophyta makes up only 2% of the phytoplankton. The Chlorophyta was dominated by *Chlorella vulgaris*, *Zygnema sterile*, and *Ankistrodesmus falcatus*, which are most abundant and occurred

throughout the study period. The Cyanophyta had *Anabaena circinalis*, *Anabaena flos-aque* and *Gomphosphaeria sp.* occurring in all the sampling stations, all year round. None of the members of Bacillariophyta and Dinophyta was present throughout the year. All the phytoplankton showed significant seasonal and spatial variation.

### Discussion

Nguru Lake is part of the Nigerian Dry Belt (Sudano – Sahelian Zone) where the scarcity of water is an important limiting factor for development (Adams and Hollis, 1989). The extent and nature of irregular and extreme ecological conditions of the savanna especially climatic variation through wind action could have a resultant effect in modifying lake ecosystems. The major physico-chemical parameters in Nguru Lake could be linked to the extent and size of the catchment area, soil characteristics, topography and vegetation covers. The water regime in this lake was mainly influenced by rainfall and discharge from its tributaries. So it is expected that any variation, whether seasonal or spatial in physico-chemical properties of Nguru lake may be influenced by climatic factors or catchment characteristics i.e. extent of human activities, and water volume fluctuations.

The marked variations in temperature observed at Nguru lake may likely be due to climatic conditions in the area, and the shallow nature of the lake. Nguru Lake being located in a Sahelian region experiences extremes in environmental temperature, this coupled with the shallowness of the lake allows for mixing. The slight seasonal variation in electrical conductivity at Nguru Lake was largely because of dilution by rains, which may most likely be due to release of ions dissolved from the previously exposed beds of the lake. This finding is consistent with the observation of Sarnelle (1992), who observed same in Lake Volta. Despite fluctuations, the water of Nguru Lake showed gradual increase in all nutrients (total phosphorus, Nitrogen and sulphates) except during the cool harmattan period. As expected, an

increase in most nutrients could be attributable to the effect of natural causes like evaporation and inflow from domestic discharges. The decrease of nutrients during the harmattan period (November – February) corresponds with the period of algal bloom. The concentration of nutrients in a water body is strongly influenced by the nature of the sediment. Wetzel (2001), states that the rate of phosphorus released into the water can double, when sediments are frequently disturbed. The values of nutrients obtained from Nguru Lake in the dry seasons were significantly higher than the respective values recorded in the rainy seasons. This could be attributed to dry season irrigation farming and concentrations of these nutrients due to evaporation. Spears *et al.*, (2007) studied seasonal partitioning of phosphorus across the sediment-water interface in Loch Leven, Scotland. They found that surface water total phosphorus concentrations were highest in late summer and lowest in early spring. In contrast, sediment total phosphorus concentrations were highest in mid-winter and lowest in late summer. In Nguru Lake, high levels of nutrients occur at the onset of the rainy season and decreased as it tends towards the dry season, in inverse proportion to plankton abundance. This agrees with fishar *et al.*, (2006) who states that nutrients were probably taken up by rapid algal and macrophyte growth in the summer giving very low nitrate and phosphate levels for the rest of the year.

The Phytoplankton was characteristically dominated by two or three species at any given time. The phytoplankton of Nguru lake were dominated by the Chlorophyta followed by Cyanophyta and Bacillariophyta with Dinophyta being the least represented. This result is similar with the findings of Abdullahi (1997) and Abubakar (2007) who worked on Hadejia-Nguru wetlands and Nguru Lake respectively. The composition of Phytoplankton in Nguru Lake was an indication that the lake is undergoing eutrophication. This is collaborated by the findings of Van Vuuren *et al.*; (2007) in Mohale dam in South Africa. However, recent trends in some sampling stations in the lake (i.e. sites 4 and 5)

indicated that the Cyanophyta was becoming more dominant in these sites, which was a clear indication of deterioration of water quality.

Observations by Odhiambo and Gichuki (1998) revealed that the Chlorophyta and the Cyanophyta dominated the algae of Lake Baringo and that the lake was in a state of gradual deterioration of water quality. These factors are applicable to Nguru Lake in the case of the present study.

### Conclusion

Different levels of anthropogenic inputs have caused wide variations in physico-chemical parameters in the various parts of Nguru Lake. There was significant seasonal variation in the abundance of the phytoplankton. The seasonal variations observed in the flora of the lake were due to availability of nutrients. The abundance of Cyanophyta in all the sampling stations is an indication that the lake is tending towards eutrophication.

### Recommendations

The government and other donor agencies should provide support for research and studies to collect analyze and synthesize information and harmonize existing policies, edicts and byelaws that conform to Integrated Water Resources Management (IWRM) principles, including establishing criteria for water – use. There should support also for the preservation of wetlands, control of erosion, restoration of degraded areas and the creation of measures to control point and non-point pollutions.

### References

**Abdullahi BA.** 1997 Biodiversity study of the aquatic fauna and flora of the Hadejia-Nguru wetlands, HNWCP, Nguru Nigeria . 79-86.

**Abubakar MM.** 2007 The phytoplankton community of a semi-and lake:Nguru lake. Kano Journal for Science Education **4(1)**,16-21.

**Adams WM.** 1993 Greening Development: Environmental issues in the third world. Routledge London .

**Adams WM, Hollis GE.** 1989 Hydrology and sustainable resource development of a Sahelian flood plain wetland. Hadejia-Nguru wetland, Conservation Project Nguru, Nigeria.

**APHA** 1985 Standard methods for examination of waters and wastewaters.16<sup>th</sup> ed. American public health authority. Washington D.C. 2112.

**Boyd CE.** 1981 Water quality in warm water fish ponds Agricultural experimental station. Gaftmaster pub. Co. Alabama USA. 359

**Crul RCM.** 1993 Monographs of the East African great Lakes. Limnology and Hydrology of Lake Victoria. Part 1. UNESCO/ IHU-IV project M-S.1 Comprehensive study of Great lakes. P68.

**Fishar MR, Thorne R, Williams WP.** 2006 Physico-chemical conditions and macro invertebrate fauna in the River Nile from Aswan to Cairo African Journal of Aquatic Science **31(2)**, 347-259.

**Hollis GE, Penson SJ, Thompson JR, Sule, AR.** 1993 Hydrology of the river basin. In Hollies *et al.*, (eds). The Hadejia-Nguru Wetlands, environment, economy and sustainable development of a Sahelian floodplain wetland IUCN Gland Switzerland.

**Immhoff KR, Alberecht D.** 1975 Oxygen dynamics in the impounded lower Ruh River Programme. Water Technology 7(314), 505-518.

**Lafiagi A.** 1997 A Report of fish biodiversity inventory in the Hadejia-Nguru Wetlands. HNWCP Nguru Nigeria.

**Odhiambo W, Gichuki J.** 1998 Seasonal dynamics of the phytoplankton community in relation to environment in Lake Baringo, Kenya. (Impact on the lake's resource management).

African Journal of Tropical Hydrobiology and Fisheries **8** 36-47.

**Prescott GW.** (1951) Algae of western Great Lake area. Cranbrook institute, Bloomfield hills, Michigan USA. 946

**Sarnelle O.** 1992 Nutrient Enrichment and grazer effect on phytoplankton in lakes. *Ecology*. **73** 551-556.

**Spears BM, Carvalho L, Paterson DM.** 2007 Phosphorus partitioning in a shallow lake: Implications for water quality management. *Water and Environment Journal*, Vol. **21** (1) 47-53pp.doi:10.1177/j.1747 – 6593. 2006. 00045. X

**Van vuuren SJ, Van derwalt N. Swanepoel, A.** 2007 Changes in algal composition and environmental variables. In the high altitude mohale Dam an important water supply reservoir to South Africa. *African Journal of Aquatic science*, **32** (3) 265. 274.

**WDI** 2006. Wetland Development Initiative. Report by the Joint Wetlands Livelihood, Hadejia, Nigeria: 3-8.

**Wetzel RG.** 2001. *Limnology, Lake and River Ecosystems (3<sup>rd</sup> edn)* Academic press. San francisco.